

PDO THREAD LIFT

Informed Consent



FULL NAME	D.O.B
<input type="text"/>	<input type="text"/>
ADDRESS	PHONE NUMBER
<input type="text"/>	<input type="text"/>

The purpose of this consent form is to provide written information regarding the risks, benefits and alternatives of the procedure named above. This material serves as a supplement to the discussion you have with your doctor/healthcare provider. It is important that you fully understand this information, so please read this document thoroughly. If you have any questions regarding the procedure, ask your doctor/healthcare professional prior to signing the consent form.

THE TREATMENT

The PDO (polydioxanone) Thread Lift and Smoothing procedure uses absorbable surgical sutures placed into the subdermal layer of the skin to initiate collagen production. The procedure can result in increased firmness and elasticity of the skin in the treated area. The nature of this cosmetic procedure may require a patient to return for numerous visits in order to achieve the desired results or to determine whether the treatment may not be completely effective at treating the particular condition.

POSSIBLE RISKS AND SIDE EFFECTS ASSOCIATED WITH PDO THREAD LIFT PROCEDURE

- Discomfort: Some discomfort may be experienced during treatment.
- Scarring: May cause scarring; sutures are inserted using a small needle, which must heal. A scar at the entry point may occur.
- Bruising, Swelling, Infection: With any minimally invasive procedure, bruising of the treated area may occur along with the potential for swelling. Bruising and swelling are very likely to occur. Infection is rare, but with any injection or incision into the skin, the possibility exists.
- Bleeding: You may experience some bleeding during the procedure. A hematoma or a small blood clot may occur and may require treatment by drainage. There is a higher risk of bleeding if you have taken any anti-inflammatory medications (Advil, Motrin, Aspirin, Ibuprofen) within the 10 days preceding the procedure.
- Damage to Deeper Structures: Deeper structures such as nerves, blood vessels and muscles may be damaged during the procedure. The potential for this to occur varies according to the location on the body the procedure is being performed. Injury to deeper structures may be temporary or permanent.
- Allergic Reaction: Allergies to tape, suture material or topical preparations have been reported. Allergic reactions may require additional treatment.
- Partial Laxity Correction: PDO Lift may not correct all your facial laxity or sagging.
- Delayed Healing: Complications may ensue as a result of smoking, using a straw, or similar motions. Smoking and similar actions are STRONGLY discouraged. Slight asymmetry, redness, visible sutures, suture breakthrough may require additional treatment or removal of the sutures.

PREGNANCY AND ALLERGIES

I am not aware that I am pregnant. I am not trying to get pregnant. I am not lactating (nursing). I do not have or have not had any major illnesses which would prohibit me from receiving dermal fillers. I certify that I do not have multiple allergies or high sensitivity to medications, including but not limited to lidocaine.